

Malay Archipelago bat not one, but two species

Genetic studies of *Myotis muricola*, otherwise known as the Wall-roosting Mouse-eared bat or Nepalese Whiskered Myotis, suggest that it consists of not one, but two distinct species.

M. muricola is widespread on the Malay Archipelago, a region with an island geography that provides natural boundaries and as a result displays some of the richest biodiversity in the world. Until now scientists had compared the shape and size of *Myotis* bats across the Archipelago and conducted only limited DNA analysis. As a result the *Myotis* family tree has remained in disarray with often contradictory lines of evidence confusing matters.

To resolve these taxonomic difficulties, a team from the Universiti Malaysia Sarawak and the Indonesian Institute of Sciences launched the first in-depth genetic study on *M. muricola* and its sister taxa, *M. mystacinus*. Writing in the *Pertanika Journal of Tropical Agricultural Science*, the team describe how they examined the bats in two regions, West and East of Wallace's line (fig. 2). Using a genetic analysis technique known as 'DNA cytochrome *b* sequencing' (fig. 3), they found that the Western and Eastern groups had a genetic distance of between 26 and 39 per cent.

Based on this high genetic distance they argue that *M. muricola* Eastern and *M. muricola* Western should be considered as two distinct species. The data also suggest that the Eastern bat started to diversify in the western region during the Pliocene (5 to 2.5 million years ago). They became fully diverged within the western region during the Pleistocene (2.5 million to 11,000 years ago) under the influence of ancient Sunda River systems that had produced gallery forest corridors which functioned as safe havens for the bats as the climate and geography around them changed.

The authors stress that as a result of these findings an official revision of the taxonomic status of *M. muricola* is urgently needed. In addition, more samples from throughout the geographic range are required to firmly establish these findings.

For more information on the research, contact

Sigit Wiantoro

Museum Zoologicum Bogoriense
Research Centre for Biology
Indonesian Institute of Sciences
Widyasatwaloka Building
Jl Raya Jakarta-Bogor Km. 46
16911 Cibinong, Indonesia.

E-mail: wiantoro@gmail.com

Tel: +(62) 2187 65056; Mobile: +(62) 8132 8891 321.

About Pertanika Journal of Tropical Agricultural Science (JTAS)

Pertanika Journal of Tropical Agricultural Science (JTAS) is published by Universiti Putra Malaysia in English and is open to authors around the world regardless of nationality. It currently publishes four times a year in February, May, August and November. Other Pertanika series include Pertanika Journal of Science & Technology (JST), and Pertanika Journal of Social Sciences & Humanities (JSSH).

JTAS aims to provide a forum for high quality research related to tropical agricultural research. Areas relevant to the scope of the journal include: agricultural biotechnology, biochemistry, biology, ecology, fisheries, forestry, food sciences, entomology, genetics, microbiology, pathology and management, physiology, plant and animal sciences, production of plants and animals of economic importance, and veterinary medicine. The journal publishes original academic articles dealing with research on issues of worldwide relevance.


Website: <http://www.pertanika.upm.edu.my/home.php>

The paper is available from this link:


[http://www.pertanika.upm.edu.my/Pertanika%20PAPERS/JTAS%20Vol.%2035%20\(2\)%20May.%202012/12%20Pg%20271-292.pdf](http://www.pertanika.upm.edu.my/Pertanika%20PAPERS/JTAS%20Vol.%2035%20(2)%20May.%202012/12%20Pg%20271-292.pdf)



1 *Myotis muricola* from Kuningan, West Java. Otherwise known as the Wall-roosting Mouse-eared bat, it is a species of vesper bat. They are typically insectivorous and nocturnal.



2 The localities of the specimens of *M. muricola* that were examined. The west of Wallace's Line region is represented by (1) Jambi (2) Krakatau Island (3) Bali Island (4) East Kalimantan (5) Southern Sarawak (6) Central Sarawak (7) Northern Sarawak (8) Tawau, Sabah (9) Fraser Hill, Pahang. The sites from the east of Wallace's Line include: (10) Lombok Island (11) Sumbawa Island (12) Sumba Island (13) Flores Island (14) Lembata Island (15) Pantar Island (16) Timor Island



3 Cytochrome b is a component of mitochondrial DNA (mtDNA) which is frequently used to determine the evolutionary relationships between organisms as it is highly variable and therefore useful for determining relationships within families and genera.

For more information about the journal, contact

The Chief Executive Editor

Journal Division
Office of the Deputy Vice Chancellor (R&I)
Tower 2, UPM-MDTC Technology Centre
Universiti Putra Malaysia
43400 Serdang, Selangor
Malaysia.

Phone: + (603) 8947 1622 | + (6) 016 217 4050

Email: ndeeps@admin.upm.edu.my